



MMWRTM

Morbidity and Mortality Weekly Report

Weekly

June 3, 2005 / Vol. 54 / No. 21

Human Exposure to Mosquito-Control Pesticides — Mississippi, North Carolina, and Virginia, 2002 and 2003

Public health officials weigh the risk for mosquito-borne diseases against the risk for human exposure to pesticides sprayed to control mosquitoes (1). Response to outbreaks of mosquito-borne diseases has focused on vector control through habitat reduction and application of pesticides that kill mosquito larvae. However, in certain situations, public health officials control adult mosquito populations by spraying ultra-low volume (ULV) (<3 fluid ounces per acre [oz/acre]) mosquito-control (MC) pesticides, such as naled, permethrin, and d-phenothrin. These ULV applications generate aerosols of fine droplets of pesticides that stay aloft and kill mosquitoes on contact while minimizing the risk for exposure to persons, wildlife, and the environment (2). This report summarizes the results of studies in Mississippi, North Carolina, and Virginia that assessed human exposure to ULV naled, permethrin, and d-phenothrin used in emergency, large-scale MC activities. The findings indicated ULV application in MC activities did not result in substantial pesticide exposure to humans; however, public health interventions should focus on the reduction of home and workplace exposure to pesticides.

FIGURE. Ultra-low volume, truck-mounted spraying for mosquito control — Mississippi, 2002



Photo/CDC

Mississippi, 2002

The 2002 West Nile virus (WNV) epidemic in Mississippi prompted an increase in MC activities, including application of ULV permethrin by truck-mounted foggers (Figure). Because of concerns about potential health effects from pesticides, the Mississippi Department of Health and CDC assessed whether MC activities increased individual urine pesticide metabolite concentrations. During September 8–19, 2002, investigators selected a geographically-random sample of 125 persons by using maps of two regions where public health officials applied MC pesticides and 67 persons from

The *MMWR* series of publications is published by the Coordinating Center for Health Information and Service, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30333.

SUGGESTED CITATION

Centers for Disease Control and Prevention. [Article Title]. *MMWR* 2005;54:[inclusive page numbers].

Centers for Disease Control and Prevention

Julie L. Gerberding, MD, MPH
Director

Dixie E. Snider, MD, MPH
Chief Science Officer

Tanja Popovic, MD, PhD
(Acting) Associate Director for Science

Coordinating Center for Health Information and Service

Blake Caldwell, MD, MPH, and Edward J. Sondik, PhD
(Acting) Directors

National Center for Health Marketing*

Steven L. Solomon, MD
(Acting) Director

Division of Scientific Communications*

Maria S. Parker
(Acting) Director

Mary Lou Lindegren, MD
(Acting) Editor, MMWR Series

Suzanne M. Hewitt, MPA
Managing Editor, MMWR Series

Douglas W. Weatherwax
(Acting) Lead Technical Writer-Editor

Stephanie M. Neitzel
Jude C. Rutledge
Writers-Editors

Lynda G. Cupell
Malbea A. LaPete
Visual Information Specialists

Kim L. Bright, MBA
Quang M. Doan, MBA
Erica R. Shaver

Information Technology Specialists

Notifiable Disease Morbidity and 122 Cities Mortality Data

Patsy A. Hall	Donna Edwards
Deborah A. Adams	Tambra McGee
Felicia J. Connor	Pearl C. Sharp
Rosaline Dhara	

* Proposed.

two control regions. Each participant completed a questionnaire describing home and occupational use of pesticides and provided a spot urine sample for analysis of pesticide metabolites 1–4 days after MC (i.e., within 5 half-lives). By using a cross-sectional design, investigators compared urine pesticide metabolite concentrations of exposed and unexposed study participants. Exposure to permethrin was verified by cross-referencing the global positioning systems location of participants with local MC spray routes. Permethrin was applied in MC regions at a concentration of 0.032 oz/acre.

Urine samples were analyzed at CDC by using tandem mass spectrometry (3). Urinary metabolite concentrations of 3-phenoxybenzoic acid (3pba), a metabolite of synthetic pyrethroid pesticides such as permethrin, did not differ significantly between MC and non-MC regions (geometric mean [GM] = 1.25 $\mu\text{g/L}$ versus 1.13 $\mu\text{g/L}$, respectively). Although 3pba concentrations did not differ between participants who used pesticides at home or at work and those who did not, participants who used pesticides on pets (n = 17) had significantly higher (p = 0.02) mean 3pba concentrations than those who did not (n = 174) (4.27 $\mu\text{g/L}$ versus 1.07 $\mu\text{g/L}$, respectively). **These findings indicated that local MC activities did not lead to increased pesticide metabolite concentrations in the urine of participants.**

North Carolina, 2003

Hurricane Isabel made landfall in North Carolina on September 18, 2003. Because of ensuing rains and flooding, mosquito populations were expected to surge. To control mosquitoes and prevent transmission of WNV and other arboviruses, the North Carolina Department of Environmental and Natural Resources (NCDENR) sprayed ULV naled and permethrin.

The North Carolina Department of Health and Human Services, NCDENR, and CDC conducted a prospective exposure assessment of ULV spraying of pesticides. Investigators recruited 90 persons from a random sample of census blocks (that accounted for the population density) marked for spraying. Participants then completed a pre-spray questionnaire about household and occupational exposure to pesticides and provided urine samples to quantify concentrations of pesticide metabolites. On September 30, aircraft in North Carolina sprayed ULV naled at 0.7 oz/acre. In addition, trucks sprayed ULV permethrin (Biomist 30+30[®]) at 0.0014 lbs/acre. Eighteen hours after aerial spraying (approximately one half-life), each participant completed a post-spray questionnaire about household and occupational exposure to pesticides and provided a second urine sample. Urine samples were analyzed at CDC by using tandem mass spectrometry (3).

Of the 90 persons recruited to participate in this exposure assessment, 75 (83%) provided pre-spray and post-spray questionnaires and urine samples. The concentrations of all pre- and post-spray pesticide metabolites measured in participant urine samples were low (Table). Dimethylphosphate (DMP), a metabolite of organophosphate pesticides such as naled, was detected in 46% of pre-spray and 49% of post-spray urine samples (limit of detection [LOD] = 0.5 µg/L). The GM 3pba concentration from post-spray urine sampled was 0.2 µg/L. Generalized estimating equations (GEE) indicated no statistically significant differences in the urine concentrations of naled and permethrin metabolites before and after spraying. Participants who ate fresh fruits or vegetables ≤3 days before completing the pre-spray (n = 58) or post-spray (n = 37) questionnaires had significantly higher urine concentrations of dimethylthiophosphate than participants who did not pre-spray (n = 16) or post-spray (n = 37) (pre-spray: 3.2 µg/L versus 1.4 µg/L; GEE p = 0.02) (post-spray: 3.3 µg/L versus 1.2 µg/L; GEE p = 0.01). Two participants who worked on farms and/or handled pesticides had significantly higher urine concentrations of nonspecific organophosphorus pesticide metabolites (e.g., dimethyldithiophosphate, diethylthiophosphate, and diethylphosphate) than participants who did not work on farms (n = 73) or handle pesticides (n = 72).

Virginia, 2003

To control mosquitoes and prevent transmission of arboviruses after Hurricane Isabel, the Virginia Department of Health (VDH) decided to spray ULV naled and d-phenothrin. VDH and CDC assessed exposure to ULV spraying of pesticides by randomly selecting 95 residents of high population-density census blocks marked for spraying. Participants then com-

pleted pre-spray questionnaires about household and occupational exposure to pesticides and provided urine samples to quantify concentrations of pesticide metabolites.

On September 30, aircraft sprayed ULV naled at 0.5 oz/acre while trucks sprayed ULV of d-phenothrin (Anvil 10+10[®]) at 0.0036 lbs/acre. Eighteen hours after spraying (approximately one half-life), each participant completed a post-spray questionnaire about household and occupational exposure to pesticides and provided a second urine sample. Urine samples were analyzed at CDC by using tandem mass spectrometry (3).

Of the 95 persons recruited for the assessment, 83 (87%) provided pre-spray and post-spray exposure questionnaires and urine samples. The concentrations of all pesticide metabolites measured in participants' urine samples were low (Table). DMP was detected in 42% of pre-spray and 48% of post-spray urine samples (LOD = 0.5 µg/L). The geometric mean 3pba concentration from post-spray urine samples was 0.6 µg/L. GEEs indicated no overall difference in the urine concentrations of naled and d-phenothrin metabolites before and after spraying.

Reported by: M Currier, MD, Univ of Mississippi Medical Center; M McNeill, MD, Mississippi Dept of Health. D Campbell, MD, North Carolina Dept of Health and Human Svcs; N Newton, PhD, North Carolina Dept of Environment and Natural Resources. JS Marr, MD, E Perry, MD, SW Berg, MD, Virginia Dept of Health. DB Barr, PhD, Div of Laboratory Sciences, GE Lubber, PhD, SM Kieszak, MA, HS Rogers, PhD, LC Backer, PhD, MG Belson, MD, C Rubin, DVM, Div of Environmental Hazards and Health Effects, National Center for Environmental Health; E Azziz-Baumgartner, MD, ZH Duprey, DVM, EIS officers, CDC.

Editorial Note: Although ULV applications of naled and synthetic pyrethroids have a low toxicity to humans, occupational

TABLE. Pre-spray and post-spray geometric mean concentrations (µg/L) of urine pesticide metabolites — North Carolina and Virginia, 2002 and 2003

Metabolite	North Carolina (n = 75)		Virginia (n = 83)		95th percentile
	Pre-spray	Post-spray	Pre-spray	Post-spray	
Dimethylphosphate*	†	†	†	†	13.0
Dimethylthiophosphate [§]	2.7	1.9	2.5	2.0	46.0
Dimethyldithiophosphate [§]	0.6	0.9	0.7	0.8	19.0
Diethylphosphate [§]	0.6	1.3	0.8	1.6	13.0
Diethylthiophosphate [§]	1.6	0.5	1.7	0.5	2.2
Diethyldithiophosphate [§]	†	†	†	†	0.9
3-Phenoxybenzoic acid [¶]	†	0.2	0.3	0.6	3.4
4-Fluoro-3-phenoxybenzoic acid	†	†	†	†	0.3
cis-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylic acid**	†	†	†	†	0.5
trans-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylic acid**	0.5	0.5	0.5	0.7	1.4
cis-3-(2,2-dibromovinyl)-2,2-dimethylcyclopropanecarboxylic acid**	†	†	†	†	0.3

* Nonspecific metabolite of naled and other organophosphate pesticides.

† Metabolite concentrations were quantitated in <50% of samples.

§ Nonspecific metabolite of organophosphate pesticides (excluding naled).

¶ Nonspecific metabolite of permethrin/d-phenothrin and other synthetic pyrethroid pesticides.

** Nonspecific metabolite of synthetic pyrethroid pesticides (excluding permethrin/d-phenothrin).

studies suggest that excessive exposure to these pesticides can cause serious health effects (4). Prolonged exposure to high concentrations of naled and synthetic pyrethroids can cause dermatitis, reactive airway disease, gastrointestinal distress, central nervous system depression, paralysis, and death (5). Exposure often results from use of these pesticides in food production, treatment of wool, wood products, and pest-control efforts; however, few studies have quantitated the level of human exposure to MC pesticides in nonoccupational settings (6).

The studies described in this report represent the first efforts to quantitate human exposure to MC pesticides during large-scale MC activities. Two of these studies used a prospective crossover design that compared urine metabolite concentrations after ULV spraying of pesticides with baseline concentrations. Use of sensitive analytic methods in these studies indicated that the urine pesticide metabolite concentrations measured were low (parts per billion). The concentration of urine metabolites in these studies are comparable with those measured in the general population (6,7). In addition, these three studies did not indicate an overall increase of pesticide metabolite concentrations in the urine of participants after spraying during MC activities. **The concentrations of naled, permethrin, and d-phenothrin during emergency ULV applications might be too low to cause important human exposure.**

In certain participants, investigators found an association between home and/or work application of pesticides and pesticide metabolite concentrations. The concentrations in participants who had histories of exposure were within the range of the general U.S. population (8). These findings are consistent with occupational studies in which prolonged exposure to pesticides through several hours of work in plant nurseries and greenhouses was associated with low but measurable concentrations of urine pesticide metabolites (9). These findings also are compatible with a prospective study that quantitated higher 3pba concentrations in the urine of pest-control operators 1 day after spraying pyrethroids (10).

The findings in this report are subject to at least three limitations. First, although naled, permethrin, and d-phenothrin remain in the environment for a short period (e.g., naled has a 1-day half-life), CDC did not conduct environmental sampling to confirm the presence of pesticide on the ground after spraying. Second, the study did not quantify the effects of synergists such as piperonyl butoxide in Anvil 10+10[®], which help increase the efficacy of synthetic pyrethroids. Finally, the use of self-reported questionnaire data limits the ability to quantify actual home or occupational pesticide exposure.

Aerial spraying with ULV naled and truck-mounted spraying with permethrin/d-phenothrin were not associated with an increase in urine pesticide metabolite concentrations among residents of these rural, suburban, and urban communities.

These findings suggest that ULV application of naled, permethrin, and d-phenothrin is safe to humans as part of integrated vector control.

The findings are noteworthy because ULV applications of pesticides that kill adult mosquitoes are an important tool in the public health response to WNV. Future studies should address the long-term safety of low-concentration exposure to naled and synthetic pyrethroid applications. In addition, public health interventions might be needed to reduce home and workplace exposure to pesticides.

References

1. Roche JP. Print media coverage of risk-risk tradeoffs associated with West Nile encephalitis and pesticide spraying. *J Urban Health* 2002; 79:482–90.
2. US Environmental Protection Agency. Health and safety specific chemicals regulatory actions. Washington, DC: Office of Pesticide Programs; 2002. Available at <http://www.epa.gov/pesticides/factsheets/pesticides4mosquitos.htm>.
3. Olsson AO, Baker SE, Nguyen JV, et al. A liquid chromatography–tandem mass spectrometry multiresidue method for quantification of specific metabolites of organophosphorus pesticides, synthetic pyrethroids, selected herbicides, and DEET in human urine. *Anal Chem* 2004;76:2453–61.
4. Edmundson WF, Davies JE. Occupational dermatitis from naled: a clinical report. *Arch Environ Health* 1967;15:89–91.
5. Mick DL, Gartin TD, Long KR. A case report: occupational exposure to the insecticide naled. *J Iowa Med Soc* 1970;LX:395–6.
6. Heudorf U, Angerer J. Metabolites of pyrethroid insecticides in urine specimens: current exposure in an urban population in Germany. *Environ Health Perspect* 2001;109:213–7.
7. CDC. Second national report on human exposure to environmental chemicals. Atlanta, GA: US Department of Health and Human Services, CDC; 2003. Available at <http://www.cdc.gov/exposurereport>.
8. Barr DB, Bravo R, Weerasekera G, et al. Concentrations of dialkyl phosphate metabolites of organophosphorus pesticides in the U.S. population. *Environ Health Perspect* 2004;112:186–200.
9. Kolmodin-Hedman B, Swensson A, Åkerblom M. Occupational exposure to some synthetic pyrethroids (permethrin and fenvalerate). *Arch Toxicol* 1982;50:27–33.
10. Leng G, Ranft U, Sugiri D, Hadnagy W, Berger-Preiss E, Idel H. Pyrethroids used indoors—biological monitoring of exposure to pyrethroids following an indoor pest control operation. *Int J Hyg Environ Health* 2003;206:85–92.

Acknowledgments

The findings in this report are based, in part, on contributions by W Rayburn, Albemarle Regional Health Svcs, Elizabeth City; J Engel, North Carolina Dept of Health and Human Svcs; M Tolliver, North Carolina Dept of Environment and Natural Resources; Z Kazzi, Office of Director, Agency for Toxic Substances and Disease Registry; K Johnson, C Sanchez, A Holmes, R Sabogal, M Patel, A Funk, C Bell, S Young, A Greiling, D Burmeister, Div of Environmental Hazards and Health Effects, C Dodson, Div of Laboratory Sciences, J Mason, E Hansen, J Shughart, Div of Emergency and Environmental Health Svcs, National Center for Environmental Health; A Hedley, Div of Health Examination Statistics, National Center for Health Statistics; G Shaughnessy, G Han, A Terranella, Epidemiology Program Office, CDC.